

Head Trauma

History

- Time of injury
- Mechanism (blunt vs. penetrating)
- Loss of consciousness
- Bleeding
- Past medical history
- Medications
- Evidence for multi-trauma

Signs and Symptoms

- Pain, swelling, bleeding
- Altered mental status
- Unconscious
- Respiratory distress / failure
- Vomiting
- Major traumatic mechanism of injury
- Seizure

Differential


- Skull fracture
- Brain injury (Concussion, Contusion, Hemorrhage or Laceration)
- Epidural hematoma
- Subdural hematoma
- Subarachnoid hemorrhage
- Spinal injury
- Abuse

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>if indicated</i>
	Obtain and Record GCS
	Supplemental oxygen Maintain SpO2 ≥ 90% Preferably ≥ 94%
	Prevent Oxygen desaturation events < 90%
	Blood Glucose Analysis Procedure
B	Maintain EtCO2 35 – 45 mmHg
A	IV / IO Procedure <i>if indicated</i>
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Multiple Trauma Protocol TB 6 <i>if indicated</i>
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>
	Spinal Motion Restriction Procedure / Protocol TB 8 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>
	Monitor and Reassess


**DO NOT ROUTINELY
HYPERVENTILATE**

**Evidence of
Brain Herniation:**
Unilateral or Bilateral Dilatation of
Pupils / Posturing


Hyperventilate to maintain
EtCO2 30 – 35 mmHg
See Pearls



Rapid Transport to appropriate destination
using
**Trauma and Burn:
EMS Triage and Destination Plan**



Trauma and Burn Protocol Section



Head Trauma

Eye Opening Response	Verbal Response	Motor Response
4 = Spontaneous	5 = Oriented	6 = Obeys commands
3 = To verbal stimuli	4 = Confused	5 = Localizes pain
2 = To pain	3 = Inappropriate words	4 = Withdraws from pain
1 = None	2 = Incoherent	3 = Flexion to pain or decorticate
	1 = None	2 = Extension to pain or decerebrate
		1 = None

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Back, Neuro**
- **GCS is a key performance measure used in the EMS Acute Trauma Care Toolkit.**
- **A single episode of hypoxia and / or hypotension can significantly increase morbidity and mortality with head injury.**
- **Hyperventilation in head injury:**
 - **Hyperventilation lowers CO₂ and causes vasoconstriction leading to increased intracranial pressure (ICP) and should not be done routinely.**
 - **Use in patient with evidence of herniation (blown pupil, decorticate / decerebrate posturing, bradycardia, decreasing GCS).**
 - **If hyperventilation is needed, ventilate at 14 – 18 / minute to maintain EtCO₂ between 30 - 35 mmHg.**
 - **Short term option only used for severe head injury typically GCS ≤ 8 or unresponsive.**
- **Do not place in Trendelenburg position as this may increase ICP and worsen blood pressure.**
- **Poorly fitted cervical collars may also increase ICP when applied too tightly.**
- **In areas with short transport times, Drug Assisted Airway protocol is not recommended for patients who are spontaneously breathing and who have oxygen saturations of ≥ 90% with supplemental oxygen including BIAD / BVM.**
- **Hypotension:**
 - **Limit IV fluids unless patient is hypotensive.**
 - **Increased intracranial pressure (ICP) may cause hypertension and bradycardia (Cushing's Response).**
 - **Usually indicates injury or shock unrelated to the head injury and should be aggressively treated.**
 - **Fluid resuscitation should be titrated to maintain at least a systolic BP of > 70 + 2 x the age in years.**
 - **Lowest blood pressure by age: < 31 days: > 60 mmHg. 31 days to 1 year: > 70 mmHg. Greater than 1 year: 70 + 2 x age in years.**
- **An important item to monitor and document is a change in the level of consciousness by serial examination.**
- **Consider Restraints if necessary for patient's and/or personnel's protection per the Restraint Procedure.**
- **Concussions:**
 - **Traumatic brain injuries involving any of a number of symptoms including confusion, LOC, vomiting, or headache.**
 - **Any prolonged confusion or mental status abnormality which does not return to normal within 15 minutes or any documented loss of consciousness should be evaluated by a physician ASAP.**
 - **EMS Providers should not make return-to-play decisions when evaluating an athlete with suspected concussion. This is outside the scope of practice.**